



Extremely Red Quasars in BOSS

Fred Hamann
University of California, Riverside

Nadia Zakamska
Nic Ross
Isabelle Paris
Rachael Alexandroff
Carolyn Villforth
Gordon Richards
Hanna Herbst
Kelly Denney
Jenny Greene
... the SDSS3-QSO team



Plausible evolution for quasars in massive high-z galaxies



Merger/ triggering event

Starburst ULIRG / SMG

Transition object blowout

Visible Quasar

Signatures of young quasars:

Red colors? Higher accretion rates? More common or more powerful outflows?
Higher star formation rates? Recent merger/interaction? Disturbed ISM/CO kinematics?

Extremely Red Quasars (ERQs) from 2 surveys:

SDSS-III/BOSS:

Sloan Digital Sky Survey-III / Baryon acoustic
Oscillation Spectroscopic Survey

DR12 (fall 2014):


Fainter & redder than SDSS-I/II (DR7)
Spectra of 297,301 quasars at $3600\text{\AA} - 1\mu\text{m}$
Quasar catalog (Paris+14,16)

WISE:

Wide-field Infrared Survey Explorer

AllWISE data release (Nov. 2013):

W1, W2, W3, W4 at 3.4, 4.6, 12, and $22\mu\text{m}$
96,000 detected in W3 at $\text{SNR} > 3$



Extremely Red Quasars (ERQs)

Ross+15

$r - W4 > 7.4$ (AB)


like Dust Obscured Galaxies (Dey+08)

At $z > 2$, particular odd line properties:


- 1) Very large emission-line REWs
- 2) Unusual “wingless” line profiles
- 3) Unusual line ratios

Hamann+16 (submitted):

- Fit CIV, NV emission lines in BOSS quasars
- Redshifts: 2.0 to 3.4
- Trends vs color (and L) across all of BOSS
- Find larger, more uniform ERQ samples





REW(CIV) depends
strongly on $i - W3$ color



Normal blue quasars:
<<1% have REW(CIV) > 150A

Extremely red quasars:
~50% have REW(CIV) > 150A



ERQs are unique with:


- Large REWs
- Wingless profiles
- Exotic line ratios
- (Outflow signatures)

Redefine “ERQs” using $i - W3 > 4.6$
($0.2 - 3.5 \mu\text{m}$ rest)

Also “core” ERQs: REW(CIV) $> 100 \text{ Å}$


→ A distinct population
with unique exotic physical conditions

Median Spectra



ERQs strongly favor large REWs,
wingless profiles, odd flux ratios

Strong relationship to i – W3 color




Several hundred
ERQs

$i-w3 > 4.6$

Peculiar Properties:
 Large REWs
 Wingless line profiles
 Large NV/CIV, NV/Ly α

And:
 Powerful Outflows




Outflow Evidence 1.:

30-65% have BAL outflow lines

2–5x greater than other BOSS quasars

Like other red quasar samples (Urrutia+09)





Outflow Evidence 2.:


Common *large* CIV blueshifts

Spikes are Ly α at $v=0$ (also OI, SiII, FeII)

~7% of ERQs have blueshifts >2500 km/s
compared to <0.1% in SDSS overall

Outflow-dominated BLRs






Outflow Evidence

3.:

VLT –
Xshooter
spectra:


[OIII] 5007



Blue quasars with REW > 150Å


ERQs with REW > 150Å

Normal [OIII], FWHM $\sim$ 500-800 km/s



Blue quasars with REW $>$ 150Å

[OIII] FWHMs & wings up to 5000 km/s
Extended ($>$ 1 kpc) quasars-driven outflows
(Zakamska+16, Hamann+ in prep.)





ERQs with REW $>$ 150Å

ERQ properties:

$L_{\text{bol}} \sim 10^{47}$ ergs/s
based on W3 mags ($\sim 3.5 \mu\text{m}$ rest)

SEDs and exotic emission lines are:

Not consistent with normal Type 1s
behind a simple dust screen



Patchy obscuration could suppress the UV
without UV reddening


Overlap with “Hot” dust obscured galaxies
(HotDOGs):

ERQs are generally less red, more uniform &
extreme AGN, similar sky densities

Polletta+07, Mateos+12, Banerli+13, Tsai+15, Assef+15

Overlap with “Hot” dust
obscured galaxies
(HotDOGs):

ERQs are generally less red,
more uniform & extreme
AGN, similar sky densities



Powerful Outflows

across scales

<1 pc to >1 kpc


BAL-like winds

Large REWs

BLR outflows

Broad blueshifted
[OIII] 5007

Closely tied to i-W3 color




Outflows enhanced by high accretion rates?
high metallicities?


Obscuration by outflowing
dusty clumpy torus?

Obscuration on galactic scales?



- Well-timed evolution between quasar properties and dusty host galaxies (seems unlikely)
- Direct interaction: Quasar outflows that shred & disperse dusty ISM clouds



Faucher-Giguere+12,13



2 ERQs with ALMA: band 6, $\sim 385 \mu\text{m}$ (rest)



Continuum: $1.57 \pm 0.026 \text{ mJy}$

$L_{\text{FIR}} \sim 3 \times 10^{12} \text{ ergs/s}$ (for Arp 220 SED)

SFR $\sim 300 \text{ Mo/yr}$

[CII](2-1):

Strong in one 100 km/s channel

Redshift problems

Summary:

- Several hundred ERQs via $i - W3$ at $2.0 < z < 3.4$:
- $L \sim 10^{47}$ ergs/s, very large REWs, odd line ratios, BAL-like outflows, BLR outflows
- Broad blueshifted [OIII] → unprecedented high-speed outflows at $> \sim 1$ kpc
- Exotic properties tied to extreme red UV- MIR color
- Not consistent with normal quasars behind a dust reddening screen
- Powerful outflows! High accretion rates? High metal abundances?
- A specific blowout phase of quasar/galaxy evolution?
- SMG-like star formation rates?
- Evolution tests: disturbed ISM/CO kinematics? higher star formation rates? (ALMA)
higher merger rates? (HST - Zakamska), ...